

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Infrared Properties of AGN in the GOODS Fields: AGN in Stellar Dominated High- z Galaxies

Jeffrey Van Duyne, C. Meg Urry, Eleni Chatzichristou, Ezequiel Treister, GOODS AGN Team

Yale University, P.O. Box 208101, New Haven, CT 06520-8101 USA

Abstract. We present analysis of spectral energy distributions (SEDs) from mid-infrared through X-ray of a sample of 420 hard X-ray selected, z -band and Spitzer/IRAC detected active galactic nuclei (AGN) and AGN candidates from the GOODS multiwavelength survey. We fit local empirical SED templates of both normal and active galaxies to the rest-frame luminosities calculated from spectroscopic (where available) and photometric redshifts. The majority of the optically stellar-dominated (with early-type galaxy fits) sources are moderately luminous ($L_{X,2-10\text{keV}} 10^{43}$ erg/s) hard X-ray sources with high X-ray hardness ratios ($HR > 0.2$), high MIR luminosities and red MIR colors in excess of a typical stellar dominated elliptical galaxy. These sources likely harbor heavily obscured (though Compton-thin) AGN. The observed ratio of obscured to unobscured AGN has an integrated mean of $\sim 3.4 : 1$ but declines with increasing redshift. This effect has been explained by Treister et al. (2004) as an observational bias triggered by the lack of spectroscopic redshifts at $R > 24$ which are predominantly higher redshift obscured sources.

The X-ray background has a far harder X-ray spectrum than the typical unobscured AGN (Mainieri et al. 2002). While a few obscured quasars ($L_{X,2-10\text{keV}} \sim 10^{45}$ erg/s) have now been observed (Norman et al. 2002, XID-S 188), that population alone does not satisfy the fraction of obscured AGN predicted by the X-ray background. Thus, we must also investigate obscured, moderate luminosity AGN out to $z \sim 1.5 - 2$, with enough dust absorption such that the host galaxy dominates the optical and NIR light with its stellar contribution. Such AGN would have significant observed hard X-ray luminosities, and mid-infrared luminosities and colors in excess of a typical elliptical galaxy indicating an obscured AGN.

With the exceptional multiwavelength coverage and depth afforded by the Great Observatories Origins Deep Survey (GOODS, Giavalisco et al. 2004), we can uncover a significant population of the obscured AGN class described above. Utilizing a sample of 420 z' -band matched hard X-ray sources in both the GOODS-N and South fields (Alexander et al. 2003), we fit AGN (obscured and unobscured) and normal galaxy (spirals and ellipticals) templates to the broadband rest-frame SEDs from U band to *Spitzer* $24\mu\text{m}/8\mu\text{m}$ (CDF-N and CDF-S fields, respectively) for sources with spectroscopic and photometric redshifts (Barger et al. 2003; Szokoly et al. 2004; Mobasher et al. 2004).

The template fits result in 82 unobscured AGN, 236 obscured AGN, and 102 other sources (normal ellipticals, starburst/starforming spirals). Approximately half of the unobscured AGN are as weak Seyfert galaxies in the local universe and have indications of a small amount of absorption (hardness ratio > 0.2 and




Figure 1. Observed obscured to unobserved ratio of AGN for GOODS fields. Vertical lines represent redshift binning. Horizontal lines are global averages at 2.8(3.6) for CDF-N(S) and 3σ errors.

($L_{8\mu\text{m}}/L_z$) > 0.5). Their SEDs fall between a heavily obscured AGN and a Seyfert 1 nucleus, indicating a partially obscured source ($N_H \sim 10^{21}$). About 65% of the obscured AGN are fit very well to a obscured AGN/Seyfert 2 template from optical to mid-IR. The remaining 35% resemble elliptical galaxies in the optical, with no obvious point source that would indicate an AGN; however, hard X-ray luminosities and IR colors reveal the obscured AGN. Normal galaxy fits with soft X-ray spectra, $L_X < 10^{42.5}$, and blue IRAC colors are *bona fide* normal galaxies. The $24\mu\text{m}$ band in MIPS shows a slight IR excess (50% luminosity excess from standard ellipticals) in 25 apparently normal elliptical galaxies in the GOODS-North field. Also, $24\mu\text{m}$ luminosities identify spiral/starburst galaxies by $[24\mu\text{m}-3.6\mu\text{m}] < -0.5$ for faint optical sources that only have photo- z . Based on IRAC/MIPS colors of such sources in the GOODS-N, we estimate 25% of the faint border-line normal galaxies with photo- z in the GOODS-S will be revealed as definite obscured AGN via $24\mu\text{m}$ observations.

From these fits we compile AGN obscured to unobserved ratios for three redshift bins (local universe, AGN evolution epoch, AGN formation epoch) for a rough estimate of ratio evolution. The observed decline in ratio towards higher redshift is expected if the intrinsic ratio is fixed over all redshifts (Treister et al. 2004), as a result of the large incompleteness in spectroscopic redshifts for sources with $R > 24$. Alternate redshifts determinations (X-ray, IR) will increase the number of both local and high- z , moderate luminosity obscured AGN. IRS spectroscopy will verify the nature of the IR bright obscured AGN candidates.

Acknowledgments. This work was supported in part by NASA grant HST-GO-09425.13-A.

References

- Alexander, D.M., Bauer, F.E., Brandt, W.N., Schneider, D.P., et al. 2003, AJ, 126, 539
- Barger, A.J., Cowie, L.L., Capak, P., Alexander, D.M. et al., 2003, AJ, 126, 632
- Giavalisco, M., Ferguson, H.C., Koekemoer, A.M., et al., 2004, ApJ, 600L, 93
- Mainieri, V., Bergeron, J., Hasinger, G., et al., 2002, A&A, 393, 425
- Mobasher, B., Idzi, R., Benitez, N., Cimatti, A. et al., 2004, ApJ, 600L, 167
- Norman, C., Hasinger, G., Giacconi, R., Gilli, R., et al., 2002, ApJ, 571, 218
- Szokoly, G.P., Bergeron, J., Hasinger, G., Lehmann, I. et al., 2004, ApJS, 155, 271
- Treister, E., Urry, C.M., Chatzichristou, E., Bauer, F. et al., 2004, ApJ, 616, 123